

Raman spectra and conformations of 2,2-dialkyl-1,3-dithio-5,6-benzocycloheptenes: coexistence of chair, boat and twist-boat forms

P.N. Sobolev, I.Kh. Shakirov and R.R. Shagidullin

A.E. Arbuzov Institute of Organic and Physical Chemistry, USSR Academy of Sciences, Kazan, 420083 (Russia)

E.N. Klimovitskii and D.Yu. Strel'nik

A.M. Butlerov Scientific Research Chemical Institute, V.I. Ul'yanov-Lenin Kazan State University, Kazan, 420008 (Russia)


(Received 29 June 1990; in final form 1 November 1991)

Abstract

Raman spectroscopy has been applied to the title compounds; dithiaketal of acetone, methyl-ethylketone, pinacolone, cyclohexanone and fluorenone. A three-component chair \rightleftharpoons boat \rightleftharpoons twist-boat conformational equilibrium has been established for all compounds studied except the pinacolone derivative, which exhibits a two-component chair \rightleftharpoons boat equilibrium. Questions concerning the symmetry of the conformations based on depolarization data are discussed.

INTRODUCTION

Seven-membered dithiaacetals (**I**) are suitable models for investigation of chair (C) \rightleftharpoons boat (B) equilibria, and the analogous acetals (**II**) provide examples of chair \rightleftharpoons twist-boat (TB) equilibria.



Experimental data for dithiaketal of acetone (**III**) [1,2], cyclohexanone (**VI**) [2,3] and fluorenone (**VII**) [4] have made it possible to determine the

Correspondence to: P.N. Sobolev, A.E. Arbuzov Institute of Organic and Physical Chemistry, USSR Academy of Sciences, Kazan, 420083, Russia; and Professor E.N. Klimovitskii, A.M. Butlerov Scientific Research Chemical Institute, V.I. Ul'yanov-Lenin Kazan State University, Kazan, 420008, Russia.